

Most of the male respondents who use tobacco products smoke cigarettes. Overall, one in five men smoke cigarettes. Among cigarette users, 8 in 10 smoked at least three cigarettes within 24 hours of prior to the interview, and nearly one-third smoked 10 or more cigarettes during the same period.

Table 3.9.1 Use of tobacco: Women

Percentage of women age 15-49 who smoke cigarettes or a pipe or use other tobacco products, according to background characteristics and maternity status, Zimbabwe 2010-11

Background characteristic	Uses tobacco				Does not use tobacco	Number of women
	Cigarettes	Pipe	Snuff	Other tobacco		
Age						
15-19	0.1	0.0	0.1	0.0	99.8	1,945
20-24	0.0	0.0	0.1	0.0	99.9	1,841
25-29	0.3	0.0	0.4	0.0	99.3	1,686
30-34	0.1	0.0	0.3	0.0	99.6	1,296
35-39	0.1	0.0	0.4	0.0	99.5	1,051
40-44	0.3	0.0	0.7	0.1	99.0	732
45-49	0.5	0.2	2.0	0.0	97.7	620
Maternity status						
Pregnant	0.0	0.0	0.4	0.1	99.5	758
Breastfeeding (not pregnant)	0.1	0.0	0.2	0.0	99.7	1,902
Neither	0.2	0.0	0.4	0.0	99.4	6,511
Residence						
Urban	0.3	0.0	0.3	0.0	99.4	3,548
Rural	0.1	0.0	0.4	0.0	99.5	5,623
Province						
Manicaland	0.0	0.0	0.4	0.0	99.6	1,227
Mashonaland Central	0.1	0.0	0.3	0.0	99.6	871
Mashonaland East	0.0	0.0	0.5	0.0	99.5	824
Mashonaland West	0.0	0.0	0.7	0.0	99.3	1,026
Matabeleland North	0.7	0.3	1.1	0.0	98.4	443
Matabeleland South	0.4	0.0	0.8	0.3	98.6	467
Midlands	0.2	0.0	0.0	0.0	99.8	1,123
Masvingo	0.0	0.0	0.1	0.0	99.9	909
Harare	0.2	0.0	0.2	0.1	99.6	1,722
Bulawayo	0.5	0.0	0.2	0.0	99.3	558
Education						
No education	0.6	0.6	1.9	0.0	97.5	212
Primary	0.2	0.0	0.6	0.1	99.2	2,568
Secondary	0.2	0.0	0.2	0.0	99.6	5,966
More than secondary	0.0	0.0	0.2	0.0	99.8	424
Wealth quintile						
Lowest	0.2	0.1	0.6	0.0	99.3	1,546
Second	0.0	0.0	0.2	0.0	99.7	1,594
Middle	0.1	0.0	0.4	0.0	99.5	1,681
Fourth	0.2	0.0	0.3	0.0	99.5	2,073
Highest	0.2	0.0	0.3	0.0	99.5	2,278
Total	0.2	0.0	0.4	0.0	99.5	9,171