


# Cambodia

## Population (in thousands)

Age	Male	Female	Total
0-29	4,695	4,578	9,273
30-44	1,126	1,256	2,382
45-59	600	777	1,377
60-69	169	274	444
70-79	65	135	200
80+	16	29	45
<b>Total</b>	<b>6,670</b>	<b>7,050</b>	<b>13,720</b>

## Population Pyramid



## Percent Smoking

Prevalence in 2005	Tobacco	Cigarette
Men	31.7	30.3
Women	12.4	10.8

## Average consumed daily by smokers

... 9

## Highlights:

- The death rate from non-communicable diseases (1070 per 100,000 population) was about 2 times larger than that for communicable diseases (566 per 100,000). Tobacco was responsible for 16% of all NCDs compared with 7% of all communicable disease related deaths.

- Within the non-communicable disease group, ischaemic heart disease accounted for 176 deaths per 100,000 population aged 30 years and over, with 20% of these deaths attributed to tobacco. In contrast, cancer of the trachea, bronchus and lung accounted for 24 deaths per 100,000 population but with 73% of these deaths attributed to tobacco.

- Within the communicable disease group, deaths attributed to tobacco accounted for 15% of all lower respiratory infection deaths and 13% of all tuberculosis deaths.

- The death rate due to tobacco in men was (326 per 100,000 men aged 30 years and over) compared with women (130 per 100,000 women aged 30 years and over). The proportion of deaths attributable to tobacco was almost 17% for men and 8% for women.

- Among those who died prematurely, almost one in every 7 deaths among those aged 30-44 and one in 7 among those aged 45-59 years were attributable to tobacco use. Of those who died in the 45-59 year age bracket, tobacco use accounted for 12.1% of those whose death was due to a lower respiratory infection, 30.7% of deaths due to the category of 'all cardiovascular diseases combined', and 71.7% of those who had died from cancer of the trachea, bronchus and lung.

## WHO estimated death rates (per 100,000) and proportion attributable to tobacco, 2004

Cause of death	Both sexes					Total for ages 30 & over
	30-44	45-59	60-69	70-79	80+	
<b>Death rate: ALL CAUSES</b>	<b>745</b>	<b>1,480</b>	<b>3,321</b>	<b>7,748</b>	<b>17,622</b>	<b>1,715</b>
Death rate attributable to tobacco	107	216	393	923	1,236	217
Proportion of deaths attributable to tobacco (%)	14	15	12	12	7	13
<b>Death rate: Communicable diseases</b>	<b>394</b>	<b>466</b>	<b>806</b>	<b>1,841</b>	<b>4,730</b>	<b>566</b>
Death rate attributable to tobacco	33	31	58	143	279	42
Proportion of deaths attributable to tobacco (%)	8	7	7	8	6	7
<b>Death rate: Tuberculosis</b>	<b>111</b>	<b>225</b>	<b>467</b>	<b>590</b>	<b>875</b>	<b>211</b>
Death rate attributable to tobacco	23	25	45	62	68	28
Proportion of deaths attributable to tobacco (%)	20	11	10	10	8	13
<b>Death rate: Lower respiratory infections</b>	<b>35</b>	<b>46</b>	<b>151</b>	<b>618</b>	<b>2,220</b>	<b>98</b>
Death rate attributable to tobacco	10	6	13	81	207	14
Proportion of deaths attributable to tobacco (%)	29	12	9	13	9	15
<b>Death rate: Noncommunicable diseases</b>	<b>276</b>	<b>940</b>	<b>2,442</b>	<b>5,775</b>	<b>12,625</b>	<b>1,070</b>
Death rate attributable to tobacco	74	185	335	780	957	175
Proportion of deaths attributable to tobacco (%)	27	20	14	14	8	16
<b>Death rate: All malignant neoplasms</b>	<b>61</b>	<b>257</b>	<b>542</b>	<b>888</b>	<b>1,194</b>	<b>219</b>
Death rate attributable to tobacco	10	46	106	200	183	41
Proportion of deaths attributable to tobacco (%)	16	18	19	23	15	19
<b>Death rate: Trachea, bronchus, lung cancers</b>	<b>5</b>	<b>25</b>	<b>68</b>	<b>130</b>	<b>126</b>	<b>24</b>
Death rate attributable to tobacco	3	18	51	100	85	17
Proportion of deaths attributable to tobacco (%)	64	72	75	77	67	73
<b>Death rate: All other malignant neoplasms</b>	<b>57</b>	<b>233</b>	<b>474</b>	<b>758</b>	<b>1,068</b>	<b>195</b>
Death rate attributable to tobacco	7	28	55	101	98	23
Proportion of deaths attributable to tobacco (%)	12	12	12	13	9	12
<b>Death rate: All Cardiovascular diseases</b>	<b>106</b>	<b>379</b>	<b>1,072</b>	<b>2,900</b>	<b>7,018</b>	<b>482</b>
Death rate attributable to tobacco	57	116	141	262	125	94
Proportion of deaths attributable to tobacco (%)	54	31	13	9	2	19
<b>Death rate: Ischaemic heart disease</b>	<b>41</b>	<b>143</b>	<b>420</b>	<b>989</b>	<b>2,330</b>	<b>176</b>
Death rate attributable to tobacco	23	42	53	81	48	35
Proportion of deaths attributable to tobacco (%)	57	30	13	8	2	20
<b>Death rate: Cerebrovascular disease</b>	<b>23</b>	<b>106</b>	<b>301</b>	<b>890</b>	<b>1,976</b>	<b>135</b>
Death rate attributable to tobacco	12	33	39	85	6	24
Proportion of deaths attributable to tobacco (%)	52	31	13	10	0	18
<b>Death rate: Other cardiovascular diseases</b>	<b>42</b>	<b>130</b>	<b>351</b>	<b>1,021</b>	<b>2,711</b>	<b>171</b>
Death rate attributable to tobacco	22	41	49	96	71	34
Proportion of deaths attributable to tobacco (%)	52	31	14	9	3	20
<b>Death rate: Respiratory diseases</b>	<b>16</b>	<b>58</b>	<b>229</b>	<b>729</b>	<b>2,080</b>	<b>103</b>
Death rate attributable to tobacco	6	19	79	293	627	37
Proportion of deaths attributable to tobacco (%)	37	33	35	40	30	35
<b>Death rate: Chronic obstructive pulmonary disease</b>	<b>3</b>	<b>21</b>	<b>111</b>	<b>378</b>	<b>909</b>	<b>45</b>
Death rate attributable to tobacco	2	14	65	242	508	28
Proportion of deaths attributable to tobacco (%)	80	65	59	64	56	62
<b>Death rate: Other respiratory diseases</b>	<b>14</b>	<b>37</b>	<b>119</b>	<b>352</b>	<b>1,170</b>	<b>58</b>
Death rate attributable to tobacco	4	6	14	51	119	9
Proportion of deaths attributable to tobacco (%)	29	15	12	14	10	15

Note 1: The smoking prevalence rates, and the death rates for ages 30 & over are not standardized and cannot be used for comparison between countries.

Note 2: For information on the sources of all data, please refer to Technical Note.

## WHO estimated death rates (per 100,000) and proportion attributable to tobacco, 2004

Cause of death	Men					Total for ages 30 & over	Women					Total for ages 30 & over
	30-44	45-59	60-69	70-79	80+		30-44	45-59	60-69	70-79	80+	
<b>Death rate: ALL CAUSES</b>	950	1,902	4,015	8,811	19,625	1,907	562	1,154	2,893	7,236	16,565	1,561
Death rate attributable to tobacco	168	344	721	1,472	1,975	326	52	117	190	658	846	130
Proportion of deaths attributable to tobacco (%)	18	18	18	17	10	17	9	10	7	9	5	8
<b>Death rate: Communicable diseases</b>	487	671	1,064	2,032	5,338	681	310	307	646	1,748	4,409	474
Death rate attributable to tobacco	49	52	113	192	341	63	19	15	24	120	246	26
Proportion of deaths attributable to tobacco (%)	10	8	11	9	6	9	6	5	4	7	6	6
<b>Death rate: Tuberculosis</b>	130	338	819	1,161	1,679	299	94	138	249	315	451	141
Death rate attributable to tobacco	34	48	102	145	159	49	12	8	9	22	20	11
Proportion of deaths attributable to tobacco (%)	26	14	12	13	9	16	13	6	4	7	4	8
<b>Death rate: Lower respiratory infections</b>	43	21	58	270	1,315	55	28	65	209	787	2,697	133
Death rate attributable to tobacco	15	4	10	46	173	13	6	7	15	98	225	15
Proportion of deaths attributable to tobacco (%)	34	19	17	17	13	24	22	10	7	12	8	11
<b>Death rate: Noncommunicable diseases</b>	331	1,107	2,841	6,619	13,997	1,096	227	811	2,196	5,368	11,901	1,048
Death rate attributable to tobacco	119	292	609	1,280	1,634	264	34	102	165	538	600	104
Proportion of deaths attributable to tobacco (%)	36	26	21	19	12	24	15	13	8	10	5	10
<b>Death rate: All malignant neoplasms</b>	51	260	644	1,091	1,238	209	71	255	479	790	1,171	226
Death rate attributable to tobacco	14	84	230	430	379	71	6	16	29	89	79	17
Proportion of deaths attributable to tobacco (%)	28	32	36	39	31	34	8	6	6	11	7	8
<b>Death rate: Trachea, bronchus, lung cancers</b>	6	38	128	251	235	36	3	14	31	71	68	14
Death rate attributable to tobacco	4	31	110	220	193	30	2	7	15	41	28	7
Proportion of deaths attributable to tobacco (%)	71	81	86	88	82	83	54	51	47	58	41	52
<b>Death rate: All other malignant neoplasms</b>	45	222	517	839	1,003	173	67	241	448	719	1,103	212
Death rate attributable to tobacco	10	53	120	210	187	41	4	9	14	48	51	10
Proportion of deaths attributable to tobacco (%)	23	24	23	25	19	23	6	4	3	7	5	5
<b>Death rate: All Cardiovascular diseases</b>	136	468	1,232	3,108	7,255	484	79	310	973	2,799	6,892	480
Death rate attributable to tobacco	95	171	215	292	70	134	24	74	95	247	154	61
Proportion of deaths attributable to tobacco (%)	70	37	17	9	1	28	30	24	10	9	2	13
<b>Death rate: Ischaemic heart disease</b>	60	198	517	1,231	2,577	200	23	100	360	872	2,200	157
Death rate attributable to tobacco	44	70	87	113	23	57	5	21	32	65	62	17
Proportion of deaths attributable to tobacco (%)	72	35	17	9	1	29	23	21	9	7	3	11
<b>Death rate: Cerebrovascular disease</b>	30	128	340	903	1,916	130	16	89	276	883	2,008	139
Death rate attributable to tobacco	16	46	56	74	17	31	8	23	28	90	-	19
Proportion of deaths attributable to tobacco (%)	55	36	17	8	1	24	46	26	10	10	-	14
<b>Death rate: Other cardiovascular diseases</b>	46	142	375	974	2,763	155	39	121	336	1,044	2,684	184
Death rate attributable to tobacco	35	55	71	105	31	46	11	30	35	92	92	25
Proportion of deaths attributable to tobacco (%)	76	39	19	11	1	30	28	24	10	9	3	13
<b>Death rate: Respiratory diseases</b>	20	76	331	1,100	2,868	122	13	44	167	551	1,664	89
Death rate attributable to tobacco	9	31	150	530	1,152	54	4	10	36	178	350	23
Proportion of deaths attributable to tobacco (%)	43	41	45	48	40	44	30	23	21	32	21	26
<b>Death rate: Chronic obstructive pulmonary disease</b>	3	31	178	654	1,575	60	2	13	69	244	558	33
Death rate attributable to tobacco	3	22	124	453	982	42	1	7	29	140	257	17
Proportion of deaths attributable to tobacco (%)	85	72	69	69	62	69	73	52	42	57	46	51
<b>Death rate: Other respiratory diseases</b>	17	45	152	446	1,293	61	11	31	98	306	1,106	56
Death rate attributable to tobacco	6	9	26	77	171	12	2	3	7	38	92	6
Proportion of deaths attributable to tobacco (%)	34	19	17	17	13	20	22	10	7	12	8	11

Note 1: The smoking prevalence rates, and the death rates for ages 30 & over are not standardized and cannot be used for comparison between countries.

Note 2: For information on the sources of all data, please refer to Technical Note.